

喇叭螺属一新种 (肺螺亚纲, 柄眼目, 虹蛹螺科)

郭云海¹ 周卫川² 罗泰昌¹

1. 贵州师范大学地理与生物科学学院 贵阳 550001
2. 福建出入境检验检疫局检验检疫技术中心 福州 350001

摘 要 在贵州省喀斯特地貌区陆生贝类多样性调查采得的标本中, 经鉴定得喇叭螺属 1 新种, 即兴义贝喇叭螺 *Boysidia (Bansonella) xingyiensis* sp. nov.。对新种的形态进行了详细描述, 与其近似种进行了对比。正、副模标本均保存于中国科学院动物研究所标本馆。

关键词 腹足纲, 肺螺亚纲, 柄眼目, 虹蛹螺科, 喇叭螺属, 新种.

中图分类号 Q959.212.2

兴义贝喇叭螺, 新种 *Boysidia (Bansonella) xingyiensis* sp. nov. (图 1~ 4)

正模标本, 壳高 2.00 mm, 壳宽 3.42 mm, 壳口高 1.57 mm, 壳口宽 1.42 mm。标本采自贵州省兴义县纳灰, 2004 年 10 月 2 日。

副模标本, 67 个。壳高 1.71~ 2.14 mm, 壳宽 2.85~ 3.57 mm, 壳口高 1.28~ 1.43 mm, 壳口宽

1.28~ 1.57 mm。标本采自贵州省兴义县纳灰, 2004 年 10 月 2 日。

形态特征 贝壳小, 壳质稍厚, 有光泽, 不透明, 呈矮圆锥形, 有 4 $\frac{1}{2}$ 螺层, 胚螺层 (1~ 2 个螺层) 缓慢增长, 略膨胀, 光滑, 体螺层特膨大, 其下部向下倾斜。在体螺层周缘上有一明显的龙骨状突起。壳面呈棕褐色, 并有无数成斜行排列的生




图 1~ 4 兴义贝喇叭螺, 新种 *Boysidia (Bansonella) xingyiensis* sp. nov., 正模 (holotype)
1. 贝壳侧面观 (lateral view of shell) 2. 壳口放大 (aperture enlarge) 3. 贝壳腹面观 (ventral view of shell)
4. 贝壳背面观 (dorsal view of shell)

长线和螺纹。壳顶钝，缝合线深。壳口呈椭圆形，与体螺层分离，呈喇叭状，扩大而外折。壳口不向下倾斜，而与体螺层下部平行，向上延长。壳口内有5枚齿状突起：1枚稍大的片状壁板；1枚大的下壁板；1枚片状上腭褶；1枚稍小的下腭褶；1枚乳状亚柱板。脐孔大，呈孔穴状，不被轴缘遮盖。

栖息环境：生活在阴暗潮湿、多石灰岩地区，常栖息于潮湿、多地衣、苔藓的石灰岩石壁上，岩石缝隙中，树叶下。

讨论 新种与皱褶贝喇叭螺 *Boysidia* (*Bensonella*) *pilicidens* (Benson, 1849) 在外形上较相似，但后者较小，(壳高 2.00 mm, 壳宽 1.5 mm)，有5个螺层，壳口内有9枚腭褶，3枚壁板，与前者有所区别。

REFERENCES (参考文献)

- Baker, H. B. 1935. Review of the anatomy of Pupillidae and related groups. In: Pilsbry, H. A. (ed.), *Manual of Conchology* (Ser. 2). 28: 191-226.
- Chen, D N and Gao, J-X 1987. *Economic Fauna Sinica of China, Terrestrial Mollusca*. Science Press, Beijing. 43-44. [陈德牛, 高家祥, 1987. 中国经济动物志, 陆生软体动物. 北京: 科学出版社. 43~44]
- Chen, D N, Wu, M and Zhang, G Q 1999. Studies on the genus

Boysidia Ancey from China with descriptions of six new species. *Acta Zootaxonomica Sinica*, 24 (2): 124-135. [陈德牛, 吴 岷, 张国庆, 1999. 中国喇叭螺属的研究及六新种记述. 动物分类学报, 24 (2): 124~135]

- Chen, D N and Zhang, W H 2002. Zoogeographical analysis of the *Boysidia* (Ancey, 1881) in China, description of a new species. (Pulmonata, Stylommatophora, Pupillidae). *Acta Zootaxonomica Sinica*, 27 (4): 690-697. [陈德牛, 张卫红, 2002. 中国喇叭螺属的动物地理分析及一新种记述. (肺螺亚纲, 柄眼目, 虹蛹螺科). 动物分类学报, 27 (4): 690~697]
- Haas, von F. 1937. Neue und Kritische Pupilliden. *Arch. Moll.*, 69 (1/2): 2-12, pl. 1-3. f. 1-34.
- Heude, P. M. 1882-1885. Notes sur les mollusques terrestres de la vallée du Fleuve Bleu. *Mem. D Hist. Nat. l'Emp. Chinois*. 12, pl. VIII figs. 7-8.
- Luo, T-C, Chen, D N and Zhang, G Q 2000. Two new species of Pupillidae from China. (Pulmonata: Stylommatophora: Pupillidae). *Acta Zootaxonomica Sinica*, 25 (2): 147-151. [罗泰昌, 陈德牛, 张国庆, 2000. 中国虹蛹螺科二新种. (肺螺亚纲, 柄眼目, 虹蛹螺科). 动物分类学报, 25 (2): 147~151]
- Pilsbry, H. A. 1916-1918. *Manual of Conchology*, (Ser. 2). 24: 1-xii, 1-380.
- Thompson, F. G. and Dance, S. P. 1983. Non-Marine Mollusks of Borneo, II. Pulmonata: Pupillidae, Clausiliidae. III. Prosobranchia: Hydrocenidae, Helicidae. *Bull. Florida State Mus. Biol. Sci.*, 29 (3): 101-152.
- Yen, T. C. 1939. Die Chinesisch Land- und Subwasser Gastropoden des Natur Museum Senckenberg. *Arch. Senckenb. Natur. Ges.*, 444: 75-76, pl. 6, f. 39-44.

ONE NEW SPECIES OF THE GENUS BOYSIDIA (PULMONATA, STYLOMMATOPHORA, PUPILLIDAE)

GUO Yur-Hai¹, ZHOU Wei-Chuan², LUO Tai-Chang¹

1. College of Geographical and Biological Science, Guizhou Normal University, Guiyang 550001, China

2. Fujian Entry-Exit Inspection & Quarantine Bureau, Fuzhou 350001, China

Abstract In the present paper, one new species of genus *Boysidia* are described. The material studied was collected by authors from Guizhou Province, China, 2 Oct. 2004. Type specimens are deposited in the Institute of Zoology, Chinese Academy of Sciences.

Boysidia (*Bansonella*) *xingyiensis* **sp. nov.** (Figs. 1-4)

Holotype. Alt. 2.00 mm, diam. 3.42 mm; alt. of aperture 1.57 mm, diam. of aperture 1.42 mm. Collected from Nahui Town, Xingyi County (25°1'N, 104°8'E), Guizhou Province, China, 2 Oct. 2004. Paratypes 67 specimens. Alt. 1.71-2.14 mm, diam. 2.85-3.57 mm; alt. of aperture 1.28-1.43 mm, diam. of aperture 1.28-1.57 mm. Collected from Nahui Town, Xingyi County (25°1'N, 104°8'E), Guizhou Province, China, 2 Oct. 2004.

Shell little, thick, luster, conic. Whorls $4\frac{1}{2}$, moderately convex, spire depressed-conical, embryo

whorls (1-2) increased slowly, smooth, the body whorl very inflated, and slants downward, with a keel-shaped at the periphery. Shell deep yellow brown in colour, and with innumerable slender and dense growth lines spiral lines, silky. Apex blunt, suture deep. Aperture elliptical, bell-formed, very expanded and reflexed, thick, the peristome separated from body whorl, and not oblique, but aperture nearly parallel to body whorl, with 5 teeth in the aperture: 1 big flake formed parietal lamella, 1 big infraparietal lamella, 1 flake formed upper palatal fold, 1 small lower palatal fold, 1 nipple-formed subcolumellar lamella. Umbilicus big, with hole formed.

This new species is bigger to *Boysidia* (*Bansonella*) *plicidens* (Benson, 1849), but differs obviously from the latter bigger size: the shell with whorls $4\frac{1}{2}$, alt. 2.00 mm, diam. 3.42 mm. The peristome separated from body whorl, and not

oblique, but aperture nearly parallel to body whorl, with 5 teeth in the aperture: 1 big flake-formed parietal lamella, 1 big infraparietal lamella, 1 flake-formed upper palatal fold, 1 small lower palatal fold,

1 nipple formed subcolumellar lamella. Umbilicus big, with hole-formed. The latter with 5 whorls (length 2.00 mm, breadth 1.5 mm) in large size, and with 5 whorls, 9 palatal folds, and 3 parietal lamellae.

Key words Gastropoda, Pulmonata, Stylommatophora, Pupillidae, *Boysidia*, new species.